

Petadolex

**60 softgels**

Item Catalog Number: 12753

Neurologist-recommended Petadolex R contains the patented extract of butterbur that supports healthy blood vessel tone in the brain, as well as normal blood flow in the brain.\*1

Petadolex is standardized to contain 15% of butterbur's key ingredients, petasins. It's also guaranteed to be pyrrolizidine alkaloid (PA) free, so you can use it with confidence. PAs are toxic compounds that can harm the liver. Studies demonstrate patented Petadolex is safe and well tolerated.

1. Lipton RB, Gobel H, Wilkes K, Mauskop A. Efficacy of petasites 50 and 75 mg for prophylaxis of migraine: results of a randomized, placebo-controlled study. *Neurology* 2002;58(S3):A472.

**Supplement Facts**

Serving Size 1 softgel

Servings Per Container 60

**Amount Per Serving**

Purple butterbur root (Petasites hybridus)

50 mg

(Petadolex R brand) root extract standardized to contain 7.5 mg petasins and to be free of pyrrolizidine alkaloids (PAs)

Other ingredients: medium chain triglycerides, gelatin, glycerin, water, and riboflavin color

Contains no sugar, salt, yeast, wheat, gluten, corn, soy, dairy products, artificial flavoring, or preservatives. All colors used are from natural sources.

**Dosage and Use**

- One softgel three times daily for the first four weeks. Thereafter, one softgel twice daily.

**Caution**

Do not use if pregnant or nursing.

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.