

Fibrinogen Resist

**30 vegetarian capsules**

Item Catalog Number: 00718

Studies indicate that excess fibrinogen is a strong predictive factor of mortality.<sup>70,71</sup> It has been suggested that plasma fibrinogen levels are closely and independently associated with the risk of silent cerebrovascular lesions.<sup>72,73</sup>

Life Extension® has determined that optimal fibrinogen levels should range between 215 and 300 milligrams per deciliter (mg/dL) of blood. Fibrinogen Resist™ provides targeted nutritional support for those seeking to maintain blood fibrinogen levels in healthy ranges. This formula offers a unique combination of enzymes and antioxidants that support cardiovascular health.

Each capsule of Fibrinogen Resist™ provides NSK-D™ nattokinase, the most bioactive form of this enzyme available. Nattokinase helps to maintain healthy fibrinolytic activity and clotting function and promotes healthy circulation and blood flow. Also included is Enzogenol®, a pine bark extract that has demonstrated antioxidant and fibrinogen-managing effects. In a study of smokers—who typically have elevated fibrinogen levels—the combination of Enzogenol® and vitamin C significantly helped normalize fibrinogen levels.<sup>74</sup>

References

Supplement Facts

Serving Size 1 vegetarian capsule

Servings Per Container 30

Amount Per Serving

Soy Natto Extract supplying 2000 fibrinolytic units of nattokinase (NSK-S D™ )	100 mg
Enzogenol® New Zealand Pine Bark ( <i>Pinus radiata</i> ) extract [std. for 83-90% proanthocyanidins (199 mg)]	240 mg
Vitamin C (as sodium ascorbate)	60 mg
Sodium (as sodium ascorbate)	8 mg

Other ingredients: vegetable cellulose, dicalcium phosphate, magnesium stearate, silica.

Contains soybeans. Contains corn and rice. This product contains NO milk, egg, fish, peanuts, crustacean shellfish, tree nuts, wheat, yeast, or gluten. Contains NO sugar, and no artificial sweeteners, flavors, colors, or preservatives.

NSK-SD™ is a trademark of Japan BiScience Laboratories.

Enzogenol® is a registered trademark of Enzo Nutraceuticals, Ltd.

Dosage and Use

- Take one capsule once or twice daily without food, or as recommended by a healthcare practitioner.
- Blood testing is suggested to assess dosage needs. Desirable blood fibrinogen levels range between 215-300 mg/dl.
- This formula has a unique combination of enzymes and antioxidants to support cardiovascular health.\*

Caution

Those who supplement with Fibrinogen Resist™ can test their blood after six weeks to assess their dosage needs. Desirable blood fibrinogen levels range between 215 mg/dL and 300 mg/dL, or as recommended by your physician, as these levels can vary between individuals. Do not use this product if you have a blood coagulation or blood pressure disorder.

Consult your health care practitioner before taking this product if you are taking any drug that affects blood coagulation or blood pressure such as heparin, warfarin (Coumadin®), diuretics, ACE inhibitors, or beta blockers.

Warnings

- Keep out of reach of children.
- Do not exceed recommended dose.
- Do not purchase if outer seal is broken or damaged.
- If you have a bad reaction to product discontinue use immediately.
- When using nutritional supplements, please consult with your physician if you are undergoing treatment for a medical condition or if you are pregnant or lactating.

All Contents Copyright © 1995-2009 Life Extension Foundation All rights reserved.

**LifeExtension**<sup>®</sup>

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.