

Seasonal Affective Disorder

REFERENCES

- Allen JM, Lam RW, et al. Depressive symptoms and family history in seasonal and nonseasonal mood disorders. *Am J Psychiatry*. 1993;150(3):443–8.
- Arendt J, Skene DJ. Melatonin as a chronobiotic. *Sleep Med Rev*. 2005;9(1):25–39.
- Birdsall TC. 5-Hydroxytryptophan: A clinically-effective serotonin precursor. *Altern Med Rev*. 1998;3(4):271–80.
- Carlsson A, Svennerholm L, et al. Seasonal and circadian monoamine variations in human brains examined post mortem. *Acta Psychiatr Scand Suppl*. 1980;280:75–85.
- Checkley SA, Murphy DG, et al. Melatonin rhythms in seasonal affective disorder. *Br J Psychiatry*. 1993;163:332–7.
- Childs PA, Rodin I, et al. Effect of fluoxetine on melatonin in patients with seasonal affective disorder and matched controls. *Br J Psychiatry*. 1995;166(2):196–8.
- Cott J, Hibbeln JR. Lack of seasonal mood change in Icelanders. *Am J Psychiatry*. 2001 Feb;158(2):328.
- Das YT, Bagchi M, et al. Safety of 5-hydroxy-L-tryptophan. *Toxicol Lett*. 2004 Apr 15;150(1):111–22. Review.
- DeBattista C, Doghramji K, et al. Adjunct modafinil for the short-term treatment of fatigue and sleepiness in patients with major depressive disorder: A preliminary double-blind, placebo-controlled study. *J Clin Psychiatry*. 2003;64(9):1057–64.
- Depue RA, Arbisi P, et al. Seasonal and mood independence of low basal prolactin secretion in premenopausal women with seasonal affective disorder. *Am J Psychiatry*. 1989;146(8):989–95.
- Depue RA, Arbisi P, et al. Seasonal independence of low prolactin concentration and high spontaneous eye blink rates in unipolar and bipolar II seasonal affective disorder. *Arch Gen Psychiatry*. 1990;47(4):356–64.
- Durlach J, Pages N, et al. Chronopathological forms of magnesium depletion with hypofunction or with hyperfunction of the biological clock. *Magnes Res*. 2002;15(3–4):263–8.
- Eastman CI, Young MA, et al. Bright light treatment of winter depression: A placebo-controlled trial. *Arch Gen Psychiatry*. 1998;55(10):883–9.
- Ford K. A seasonal depression: Management of seasonal affective disorder. *Prof Nurse*. 1992;8(2):94–8.
- Ghadirian AM, Murphy BE, et al. Efficacy of light versus tryptophan therapy in seasonal affective disorder. *J Affect Disord*. 1998;50(1):23–7.
- Gross F, Gysin F. [Phototherapy in psychiatry: clinical update and review of indications]. *Encephale*. 1996;22(2):143–8.
- Guilleminault C, Cathala JP, et al. Effects of 5-hydroxytryptophan on sleep of a patient with a brain-stem lesion. *Electroencephalogr Clin Neurophysiol*. 1973;34(2):177–84.
- Hebert M, Beattie CW, et al. Electroretinography in patients with winter seasonal affective disorder. *Psychiatry Res*. 2004;127(1–2):27–34.
- Hilger E, Willeit M, et al. Reboxetine in seasonal affective disorder: An open trial. *Eur Neuropsychopharmacol*. 2001;11(1):1–5.
- Hirota T, Fukada Y. Resetting mechanism of central and peripheral circadian clocks in mammals. *Zoolog Sci*. 2004;21(4):359–68.

Hoekstra R, Fekkes D, et al. Effect of light therapy on bipterin, neopterin and tryptophan in patients with seasonal affective disorder. *Psychiatry Res.* 2003;120(1):37–42.

Hvas AM, Juul S, et al. Vitamin B6 level is associated with symptoms of depression. *Psychother Psychosom.* 2004;73(6):340–3.

Ising H, Bertschat F, et al. Measurement of free magnesium in blood, serum and plasma with an ion-sensitive electrode. *Eur J Clin Chem Clin Biochem.* 1995;33(6):365–71.

Jacobsen FM, Sack DA, et al. Neuroendocrine response to 5-hydroxytryptophan in seasonal affective disorder. *Arch Gen Psychiatry.* 1987 Dec;44(12):1086–91.

Jepson TL, Ernst ME, et al. Current perspectives on the management of seasonal affective disorder. *J Am Pharm Assoc (Wash).* 1999;39(6):822–9.

Kasper S. Treatment of seasonal affective disorder (SAD) with hypericum extract. *Pharmacopsychiatry.* 1997;30(Suppl 2): 89–93.

Kasper S, el Giamal N, et al. Reboxetine: The first selective noradrenaline re-uptake inhibitor. *Expert Opin Pharmacother.* 2000;1(4):771–82.

Lam RW, Levitan RD. Pathophysiology of seasonal affective disorder: A review. *J Psychiatry Neurosci.* 2000a;25(5):469–80.

Lam RW, Bowering TA, et al. Effects of rapid tryptophan depletion in patients with seasonal affective disorder in natural summer remission. *Psychol Med.* 2000b;30(1):79–87.

Lam RW, Gorman CP, et al. Multicenter, placebo-controlled study of fluoxetine in seasonal affective disorder. *Am J Psychiatry.* 1995;152(12):1765–70.

Lam RW, Levitan RD, et al. L-tryptophan augmentation of light therapy in patients with seasonal affective disorder. *Can J Psychiatry.* 1997;42(3):303–6.

Lam RW, Zis AP, et al. Effects of rapid tryptophan depletion in patients with seasonal affective disorder in remission after light therapy. *Arch Gen Psychiatry.* 1996;53(1):41–4.

Lambert GW, Reid C, et al. Effect of sunlight and season on serotonin turnover in the brain. *Lancet.* 2002;360(9348):1840–2.

Lansdowne AT, Provost SC. Vitamin D3 enhances mood in healthy subjects during winter. *Psychopharmacology (Berl).* 1998;135(4):319–23.

Leu SJ, Shiah IS, et al. Immune-inflammatory markers in patients with seasonal affective disorder: Effects of light therapy. *J Affect Disord.* 2001;63(1–3):27–34.

Lewy AJ, Bauer VK, et al. Melatonin treatment of winter depression: A pilot study. *Psychiatry Res.* 1998a;77(1):57–61.

Lewy AJ, Bauer VK, et al. Morning vs evening light treatment of patients with winter depression. *Arch Gen Psychiatry.* 1998b;55(10):890–6.

Lewy AJ, Sack RL, et al. Antidepressant and circadian phase-shifting effects of light. *Science.* 1987a;235(4786):352–4.

Lewy AJ, Sack RL, et al. The phase shift hypothesis for bright light's therapeutic mechanism of action: Theoretical considerations and experimental evidence. *Psychopharmacol Bull.* 1987b;23(3):349–53.

Lundt L. Modafinil treatment in patients with seasonal affective disorder/winter depression: An open-label pilot study. *J Affect Disord.* 2004;81(2):173–8.

Macchi MM, Bruce JN. Human pineal physiology and functional significance of melatonin. *Front Neuroendocrinol.* 2004;25(3–4):177–95.

Madden PA, Heath AC, et al. Seasonal changes in mood and behavior: The role of genetic factors. *Arch Gen Psychiatry.* 1996;53(1):47–55.

- Magnusson A, Axelsson J, et al. Lack of seasonal mood change in the Icelandic population: Results of a cross-sectional study. *Am J Psychiatry*. 2000 Feb;157(2):234–8.
- Magnusson A, Boivin D. Seasonal affective disorder: An overview. *Chronobiol Int*. 2003;20(2):189–207.
- Magnusson A, Partonen T. The diagnosis, symptomatology, and epidemiology of seasonal affective disorder. *CNS Spectr*. 2005 Aug;10(8):625–34;quiz 1–14. Review.
- Maini RN, Taylor PC, et al. Insights into the pathogenesis of rheumatoid arthritis from application of anti-TNF therapy. *Nihon Rinsho Meneki Gakkai Kaishi*. 2000;23(6):487–9.
- Martin TG. Serotonin syndrome. *Ann Emerg Med*. 1996 Nov;28(5):520-6. Review.
- Martinez B, Kasper S, et al. Hypericum in the treatment of seasonal affective disorders. *J Geriatr Psychiatry Neurol*. 1994;7(Suppl 1):S29-S33.
- McGeer PL, McGeer EG. The inflammatory response system of brain: implications for therapy of Alzheimer and other neurodegenerative diseases. *Brain Res Brain Res Rev*. 1995 Sep;21(2):195-218. Review.
- McGrath-Hanna NK, Greene DM, et al. Diet and mental health in the Arctic: Is diet an important risk factor for mental health in circumpolar peoples?—a review. *Int J Circumpolar Health*. 2003 Sep;62(3):228–41. Review.
- McKee T, McKee J. Nitrogen metabolism I: Synthesis. In Peterson K, Horn M, eds. *Biochemistry—An Introduction*. 2nd ed. New York, NY: McGraw-Hill; 1999:359–408.
- Menza MA, Kaufman KR, et al. Modafinil augmentation of antidepressant treatment in depression. *J Clin Psychiatry*. 2000;61(5):378–81.
- Moscovitch A, Blashko CA, et al. A placebo-controlled study of sertraline in the treatment of outpatients with seasonal affective disorder. *Psychopharmacology (Berl)*. 2004;171(4):390–7.
- Nangia M, Syed W, et al. Efficacy and safety of St. John's wort for the treatment of major depression. *Public Health Nutr*. 2000;3(4A):487–94.
- Partonen T, Vakkuri O, et al. Effects of bright light on sleepiness, melatonin, and 25-hydroxyvitamin D(3) in winter seasonal affective disorder. *Biol Psychiatry*. 1996;39(10):865–72.
- Pjrek E, Winkler D, et al. Bright light therapy in seasonal affective disorder—does it suffice? *Eur Neuropsychopharmacol*. 2004;14(4):347–51.
- Potter W, Hollister L. Antidepressant agents. In Katzung B, ed. *Basic & Clinical Pharmacology*. 8th ed. New York, NY: McGraw-Hill; 2001:498–511.
- Rohr UD, Herold J. Melatonin deficiencies in women. *Maturitas*. 2002;41(Suppl 1):S85–104.
- Rosen LN, Targum SD, et al. Prevalence of seasonal affective disorder at four latitudes. *Psychiatry Res*. 1990 Feb;31(2):131-44.
- Rosenthal NE, Sack DA, et al. Seasonal affective disorder: A description of the syndrome and preliminary findings with light therapy. *Arch Gen Psychiatry*. 1984;41(1):72–80.
- Scammell TE, Estabrooke IV, et al. Hypothalamic arousal regions are activated during modafinil-induced wakefulness. *J Neurosci*. 2000;20(22):8620–28.
- Schlager DS. Early-morning administration of short-acting beta blockers for treatment of winter depression. *Am J Psychiatry*. 1994;151(9):1383–5.
- Schwartz PJ, Murphy DL, et al. Effects of meta-chlorophenylpiperazine infusions in patients with seasonal affective disorder and healthy control subjects. Diurnal responses and nocturnal regulatory mechanisms. *Arch Gen Psychiatry*. 1997;54(4):375–85.
- Sher L. Genetic studies of seasonal affective disorder and seasonality. *Compr Psychiatry*. 2001;42(2):105–10.

Soulairac A, Lambinet H. [Effect of 5-hydroxytryptophan, a serotonin precursor, on sleep disorders]. *Ann Med Psychol (Paris)*. 1977;1(5):792–8.

Stastny J, Konstantinidis A, et al. Effects of tryptophan depletion and catecholamine depletion on immune parameters in patients with seasonal affective disorder in remission with light therapy. *Biol Psychiatry*. 2003;53(4):332–7.

Terman JS, Terman M. Photopic and scotopic light detection in patients with seasonal affective disorder and control subjects. *Biol Psychiatry*. 1999;46(12):1642–8.

Terman JS, Terman M, et al. Circadian time of morning light administration and therapeutic response in winter depression. *Arch Gen Psychiatry*. 2001;58(1):69–75.

Thompson C, Isaacs G. Seasonal affective disorder—a British sample: Symptomatology in relation to mode of referral and diagnostic subtype. *J Affect Disord*. 1988;14(1):1–11.

Thornton AE, Raz N. Memory impairment in multiple sclerosis: A quantitative review. *Neuropsychology*. 1997;11(3):357–66.

Touitou Y, Touitou C, et al. Serum magnesium circadian rhythm in human adults with respect to age, sex and mental status. *Clin Chim Acta*. 1978;87(1):35–41.

Vorbach EU, Arnoldt KH, Hubner WD. Efficacy and tolerability of St. John's wort extract LI 160 versus imipramine in patients with severe depressive episodes according to ICD-10. *Pharmacopsychiatry*. 1997 Sep;30 Suppl 2:81-5.

White DM, Lewy AJ, et al. Is winter depression a bipolar disorder? *Compr Psychiatry*. 1990;31(3):196–204.

Winton F, Corn T, et al. Effects of light treatment upon mood and melatonin in patients with seasonal affective disorder. *Psychol Med*. 1989;19(3):585–90.

Wirz-Justice A, Bucheli C, et al. Light treatment of seasonal affective disorder in Switzerland. *Acta Psychiatr Scand*. 1986;74(2):193–204.

Wyatt RJ, Zarcone V, et al. Effects of 5-hydroxytryptophan on the sleep of normal human subjects. *Electroencephalogr Clin Neurophysiol*. 1971 Jun;30(6):505-9.

All Contents Copyright © 1995-2009 Life Extension Foundation All rights reserved.

LifeExtension®

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.