

## Osteoarthritis

### REFERENCES

- Adcocks C, Collin P, et al. Catechins from green tea (*Camellia sinensis*) inhibit bovine and human cartilage proteoglycan and type II collagen degradation in vitro. *J Nutr*. 2002 Mar;132(3):341-6.
- Ahmed S, Rahman A, et al. Green tea polyphenol epigallocatechin-3-gallate inhibits the IL-1 beta-induced activity and expression of cyclooxygenase-2 and nitric oxide synthase-2 in human chondrocytes. *Free Radic Biol Med*. 2002 Oct 15;33(8):1097-105.
- Ahmed S, Wang N, et al. Green tea polyphenol epigallocatechin-3-gallate (EGCG) differentially inhibits interleukin-1 beta-induced expression of matrix metalloproteinase-1 and -13 in human chondrocytes. *J Pharmacol Exp Ther*. 2004 Feb;308(2):767-73.
- Altman RD, Marcussen KC. Effects of a ginger extract on knee pain in patients with osteoarthritis. *Arthritis Rheum*. 2001 Dec;44(11):2531-8.
- Appelboom T, Schuermans J, et al. Symptoms modifying effect of avocado/soybean unsaponifiables (ASU) in knee osteoarthritis: a double blind, prospective, placebo-controlled study. *Scand J Rheumatol*. 2001;30(4):242-7.
- Banerjee M, Tripathi LM, et al. Modulation of inflammatory mediators by ibuprofen and curcumin treatment during chronic inflammation in rat. *Immunopharmacol Immunotoxicol*. 2003 May;25(2):213-24.
- Barcelo HA, Wiemeyer JC, et al. Experimental osteoarthritis and its course when treated with S-adenosyl-L-methionine [in Spanish]. *Rev Clin Esp*. 1990 Jun;187(2):74-8.
- Blotman F, Maheu E, et al. Efficacy and safety of avocado/soybean unsaponifiables in the treatment of symptomatic osteoarthritis of the knee and hip: a prospective, multicenter, three-month, randomized, double-blind, placebo-controlled trial. *Rev Rhum Engl Ed*. 1997 Dec;64(12):825-34.
- Bucsi L, Poor G. Efficacy and tolerability of oral chondroitin sulfate as a symptomatic slow-acting drug for osteoarthritis (SYSADOA) in the treatment of knee osteoarthritis. *Osteoarthritis Cartilage*. 1998 May;6(suppl A):31-6.
- Fronzoza CG, Sohrabi A, et al. An in vitro screening assay for inhibitors of proinflammatory mediators in herbal extracts using human synovial cell cultures. *In Vitro Cell Dev Biol Anim*. 2004 Mar-Apr;40(3-4):95-101.
- Glorioso S, Todesco S, et al. Double-blind multicentre study of the activity of S-adenosylmethionine in hip and knee osteoarthritis. *Int J Clin Pharmacol Res*. 1985;5(1):39-49.
- Henrotin YE, Labasse AH, et al. Effects of three avocado/soybean unsaponifiable mixtures on metalloproteinases, cytokines and prostaglandin E2 production by human articular chondrocytes. *Clin Rheumatol*. 1998;17(1):31-9.
- Huang MH, Yang RC, et al. Preliminary results of integrated therapy for patients with knee osteoarthritis. *Arthritis Rheum*. 2005 Dec 15;53(6):812-20.
- Ishiwa J, Sato T, et al. A citrus flavonoid, nobiletin, suppresses production and gene expression of matrix metalloproteinase 9/gelatinase B in rabbit synovial fibroblasts. *J Rheumatol*. 2000 Jan;27(1):20-5.
- Joe B, Lokesh BR. Effect of curcumin and capsaicin on arachidonic acid metabolism and lysosomal enzyme secretion by rat peritoneal macrophages. *Lipids*. 1997 Dec;32(11):1173-80.
- Kalbhen DA, Jansen G. Pharmacologic studies on the antidegenerative effect of ademetionine in experimental arthritis in animals [in German]. *Arzneimittelforschung*. 1990 Sep;40(9):1017-21.
- Kasper DL, Braunwald E, et al, eds. *Harrison's Principles of Internal Medicine*. 16th ed. New York, NY. McGraw-Hill; 2004.

Kim LS, Axelrod LJ, et al. Efficacy of MSM in osteoarthritis pain of the knee: a pilot clinical trial. *Osteoarthritis Cartilage*. 2006 Mar;14(3):286-94.

Lawrence RM. Methylsulfonylmethane (MSM): a double-blind study of its use in degenerative arthritis. *Int J Anti-Aging Med*. 1998;1:50.

Lieber CS, Packer L. S-Adenosylmethionine: molecular, biological, and clinical aspects—an introduction. *Am J Clin Nutr*. 2002 Dec;76(5):1148S-50S.

Lin N, Sato T, et al. Novel anti-inflammatory actions of nobiletin, a citrus polymethoxy flavonoid, on human synovial fibroblasts and mouse macrophages. *Biochem Pharmacol*. 2003 Jun 15;65(12):2065-71.

Manthey JA, Grohmann K, et al. Biological properties of citrus flavonoids pertaining to cancer and inflammation. *Curr Med Chem*. 2001 Feb;8(2):135-53.

Mauviel A, Loyau G, et al. Effect of unsaponifiable extracts of avocado and soybean (Piascledine) on the collagenolytic action of cultures of human rheumatoid synoviocytes and rabbit articular chondrocytes treated with interleukin-1 [in French]. *Rev Rhum Mal Osteoartic*. 1991 Apr;58(4):241-5.

Miggiano GA, Gagliardi L. Diet, nutrition and rheumatoid arthritis [in Italian]. *Clin Ter*. 2005 May-Jun;156(3):115-23.

Mori TA, Beilin LJ. Omega-3 fatty acids and inflammation. *Curr Atheroscler Rep*. 2004 Nov;6(6):461-7.

Murakami A, Nakamura Y, et al. Inhibitory effect of citrus nobiletin on phorbol ester-induced skin inflammation, oxidative stress, and tumor promotion in mice. *Cancer Res*. 2000a Sep 15;60(18):5059-66.

Murakami A, Nakamura Y, et al. Suppressive effects of citrus fruits on free radical generation and nobiletin, an anti-inflammatory polymethoxyflavonoid. *Biofactors*. 2000b;12(1-4):187-92.

Murav'ev I, Venikova MS, et al. Effect of dimethyl sulfoxide and dimethyl sulfone on a destructive process in the joints of mice with spontaneous arthritis [in Russian]. *Patol Fiziol Eksp Ter*. 1991 Mar-Apr;(2):37-9.

Neustadt D, Caldwell J, et al. Clinical effects of intraarticular injection of high molecular weight hyaluronan (Orthovisc) in osteoarthritis of the knee: a randomized, controlled, multicenter trial. *J Rheumatol*. 2005 Oct;32(10):1928-36.

Obertreis B, Ruttkowski T, et al. Ex-vivo in-vitro inhibition of lipopolysaccharide stimulated tumor necrosis factor-alpha and interleukin-1 beta secretion in human whole blood by extractum urticae dioicae foliorum. *Arzneimittelforschung*. 1996 Apr;46(4):389-94.

O'Leary KA, de Pascual-Tereasa S, et al. Effect of flavonoids and vitamin E on cyclooxygenase-2 (COX-2) transcription. *Mutat Res*. 2004 Jul;551(1-2):245-54.

Plummer SM, Holloway KA, et al. Inhibition of cyclo-oxygenase 2 expression in colon cells by the chemopreventive agent curcumin involves inhibition of NF-kappaB activation via the NIK/IKK signalling complex. *Oncogene*. 1999 Oct 28;18(44):6013-20.

Podoprigrorova VG, Bobylev AA, et al. Peculiarities of free radical and antioxidant statute in patients with systemic lupus erythematosus and rheumatoid arthritis [in Russian]. *Biomed Khim*. 2005 Sep-Oct;51(5):536-41.

Qiu GX, Gao SN, et al. Efficacy and safety of glucosamine sulfate versus ibuprofen in patients with knee osteoarthritis. *Arzneimittelforschung*. 1998 Jun;48(5):469-74.

Regan E, Flannelly J, et al. Extracellular superoxide dismutase and oxidant damage in osteoarthritis. *Arthritis Rheum*. 2005 Nov;52(11):3479-91.

Rizzo R, Grandolfo M, et al. Calcium, sulfur, and zinc distribution in normal and arthritic articular equine cartilage: a synchrotron radiation-induced X-ray emission (SRIXE) study. *J Exp Zool*. 1995 Sep 1;273(1):82-6.

Salk R, Chang T, et al. Viscosupplementation (hyaluronans) in the treatment of ankle osteoarthritis. *Clin Podiatr Med Surg North Am*. 2005 Oct;22(4):585-97, vii.

Schulze-Tanzil G, de SP, et al. Effects of the antirheumatic remedy hox alpha—a new stinging nettle leaf extract—on matrix metalloproteinases in human chondrocytes in vitro. *Histol Histopathol.* 2002 Apr;17(2):477-85.

Singh R, Ahmed S, et al. Epigallocatechin-3-gallate selectively inhibits interleukin-1beta-induced activation of mitogen activated protein kinase subgroup c-Jun N-terminal kinase in human osteoarthritis chondrocytes. *J Orthop Res.* 2003 Jan;21(1):102-9.

Soeken KL, Lee WL, et al. Safety and efficacy of S-adenosylmethionine (SAME) for osteoarthritis. *J Fam Pract.* 2002 Jun;51(5):425-30.

Srivastava KC, Mustafa T. Ginger (*Zingiber officinale*) in rheumatism and musculoskeletal disorders. *Med Hypotheses.* 1992 Dec;39(4):342-8.

Tanaka S, Sato T, et al. Prevention of UVB-induced photoinflammation and photoaging by a polymethoxy flavonoid, nobiletin, in human keratinocytes in vivo and in vitro. *Biochem Pharmacol.* 2004 Aug 1;68(3):433-9.

Theiler R, Bruhlmann P. Overall tolerability and analgesic activity of intra-articular sodium hyaluronate in the treatment of knee osteoarthritis. *Curr Med Res Opin.* 2005 Nov;21(11):1727-33.

Usha PR, Naidu MU. Randomised, double-blind, parallel, placebo-controlled study of oral glucosamine, methylsulfonylmethane and their combination in osteoarthritis. *Clin Drug Invest.* 2004 Jun;24(6):353-63.

Verbruggen G, Goemaere S, et al. Chondroitin sulfate: S/DMOAD (structure/disease modifying anti-osteoarthritis drug) in the treatment of finger joint OA. *Osteoarthritis Cartilage.* 1998 May;6(suppl A):37-8.

Vetter G. Double-blind comparative clinical trial with S-adenosylmethionine and indomethacin in the treatment of osteoarthritis. *Am J Med.* 1987 Nov 20;83(5A):78-80.

Zak A, Tyrzicka E, et al. Pathophysiology of and clinical significance of polyunsaturated fatty acids n-3 family [in Czech]. *Cas Lek Cesk.* 2005;144(suppl 1):6-18.

All Contents Copyright © 1995-2009 Life Extension Foundation All rights reserved.

**LifeExtension**<sup>®</sup>

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.