

Hemochromatosis

REFERENCES

- Borgaonkar MR. Hemochromatosis: More common than you think. *Can Fam Physician*. 2003 Jan;49:36–43.
- Britton RS, Leicester KL, et al. Iron toxicity and chelation therapy. *Int Journal Hematology*. 2002 Oct;76(3):219–28.
- Chawla RK, Bonkovsky HL, et al. Biochemistry and pharmacology of S-adenosyl-L-methionine and rationale for its use in liver disease. *Drugs*. 1990;40 Suppl 3:98-110.
- Crowe S. Amlodipine decreases iron uptake and oxygen free radical production in the heart of chronically iron overloaded mice. *Biology Res Nursing*. 2002 Apr;3(4):189–97.
- Hallberg L. Does calcium interfere with iron absorption? *Am J Clin Nutr*. 1998 Jul;68(1):3–4.
- Kaltwasser JP, Werner E, et al. Clinical trial on the effect of regular tea drinking on iron accumulation in genetic haemochromatosis. *Gut*. 1998;43(5):699–704.
- Kasper DL, Braunwald E, et al. *Harrison's Principles of Internal Medicine*. 16th ed. New York: McGraw-Hill; 2005: 2298–303.
- Kidd P. Glutathione: Systemic protectant against oxidative and free radical damage. *Alt Med Review*. 2005.
- Murray M. Iron. In: Murphy MT, ed. *Encyclopedia of Nutritional Supplements*. Rocklin, Calif: Prima Publishing; 1996:208.
- Young IS, Trouton TG, et al. Antioxidant status and lipid peroxidation in hereditary haemochromatosis. *Free Radic Biol Med*. 1994 Mar;16(3):393–7.

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.