

Rheumatoid Arthritis

REFERENCES

- Adam O, Beringer C, et al. Anti-inflammatory effects of a low arachidonic acid diet and fish oil in patients with rheumatoid arthritis. *Rheumatol Int.* 2003 Jan;23(1):27–36.
- Adcocks C, Collin P, et al. Catechins from green tea (*Camellia sinensis*) inhibit bovine and human cartilage proteoglycan and type II collagen degradation in vitro. *J Nutr.* 2002 Mar;132(3):341–6.
- Ahmed S, Rahman A, et al. Green tea polyphenol epigallocatechin-3-gallate inhibits the interleukin-1 beta-induced activity and expression of cyclooxygenase-2 and nitric oxide synthase-2 in human chondrocytes. *Free Radic Biol Med.* 2002 Oct 15;33(8):1097–105.
- Ahmed S, Wang N, et al. Green tea polyphenol epigallocatechin-3-gallate (EGCG) differentially inhibits interleukin-1 beta-induced expression of matrix metalloproteinase-1 and -13 in human chondrocytes. *J Pharmacol Exp Ther.* 2004 Feb;308(2):767–73.
- Ammon HP. [Boswellic acids (components of frankincense) as the active principle in treatment of chronic inflammatory diseases]. *Wien Med Wochenschr.* 2002;152(15-16):373–8.
- Banerjee M, Tripathi LM, et al. Modulation of inflammatory mediators by ibuprofen and curcumin treatment during chronic inflammation in rat. *Immunopharmacol Immunotoxicol.* 2003 Jun;25(2):213–24.
- Bertolini DR, Nedwin GE, et al. Stimulation of bone resorption and inhibition of bone formation in vitro by human tumour necrosis factors. *Nature.* 1986 Feb 6;319(6053):516–8.
- Cai WC, Wang H, et al. [Determination of oxidation-reduction level in patients with rheumatoid arthritis]. *Di Yi Jun Yi Da Xue Xue Bao.* 2005 Jun;25(6):749–50. Chinese.
- Capellino S, Montagna P, et al. [Effects of estrogen peripheral metabolism in rheumatoid arthritis.]. *Reumatismo.* 2005 Apr-Jun;57(2):78–82.
- Cerhan JR, Saag KG, et al. Antioxidant micronutrients and risk of rheumatoid arthritis in a cohort of older women. *Am J Epidemiol.* 2003 Feb 15;157(4):345–54.
- Fronzoza CG, Sohrabi A, et al. An in vitro screening assay for inhibitors of proinflammatory mediators in herbal extracts using human synovioyte cultures. *In Vitro Cell Dev Biol Anim.* 2004 Mar;40(3-4):95–101.
- Goggs R, Vaughan-Thomas A, et al. Nutraceutical therapies for degenerative joint diseases: A critical review. *Crit Rev Food Sci Nutr.* 2005;45(3):145–64. Review.
- Gutierrez S, Palacios I, et al. SAME restores the changes in the proliferation and in the synthesis of fibronectin and proteoglycans induced by tumour necrosis factor alpha on cultured rabbit synovial cells. *Br J Rheumatol.* 1997 Feb;36(1):27–31.
- Hua J, Suguro S, et al. Preventive actions of a high dose of glucosamine on adjuvant arthritis in rats. *Inflamm Res.* 2005 Mar;54(3):127–32.
- Imrich R, Rovensky J, et al. Low levels of dehydroepiandrosterone sulphate in plasma, and reduced sympathoadrenal response to hypoglycaemia in premenopausal women with rheumatoid arthritis. *Ann Rheum Dis.* 2005 Feb;64(2):202–6.
- Ishiwa J, Sato T, et al. A citrus flavonoid, nobiletin, suppresses production and gene expression of matrix metalloproteinase 9/gelatinase B in rabbit synovial fibroblasts. *J Rheumatol.* 2000 Jan;27(1):20–5.
- Joe B, Lokesh BR. Effect of curcumin and capsaicin on arachidonic acid metabolism and lysosomal enzyme secretion by rat peritoneal macrophages. *Lipids.* 1997 Dec;32(11):1173–80.

Kamanli A, Naziroglu M, et al. Plasma lipid peroxidation and antioxidant levels in patients with rheumatoid arthritis. *Cell Biochem Funct.* 2004 Jan-Feb;22(1):53–7.

Kasper DL, Braunwald E, et al. *Harrison's Principles of Internal Medicine.* 16th ed. New York: McGraw-Hill Medical Publishing Division; 2005.

Kast RE. Borage oil reduction of rheumatoid arthritis activity may be mediated by increased cAMP that suppresses tumor necrosis factor-alpha. *Int Immunopharmacol.* 2001 Dec;1(12):2197–9.

Kremer JM, Lawrence DA, et al. Effects of high-dose fish oil on rheumatoid arthritis after stopping NSAIDs: Clinical and immune correlates. *Arthritis Rheum.* 1995 Aug;38(8):1107–14.

Lieber CS, Packer L. S-Adenosylmethionine: Molecular, biological, and clinical aspects—an introduction. *Am J Clin Nutr.* 2002 Dec;76(5):1148S–1150S.

Lin N, Sato T, et al. Novel anti-inflammatory actions of nobiletin, a citrus polymethoxy flavonoid, on human synovial fibroblasts and mouse macrophages. *Biochem Pharmacol.* 2003 Jun 15;65(12):2065–71.

Manthey JA, Grohmann K, et al. Biological properties of citrus flavonoids pertaining to cancer and inflammation. *Curr Med Chem.* 2001 Feb;8(2):135–53.

Murakami A, Nakamura Y, et al. Inhibitory effect of citrus nobiletin on phorbol ester-induced skin inflammation, oxidative stress, and tumor production in mice. *Cancer Res.* 2000b Sep 15;60(18):5059–66.

Murakami A, Nakamura Y, et al. Suppressive effects of citrus fruits on free radical generation and nobiletin, an anti-inflammatory polymethoxyflavonoid. *Biofactors.* 2000a;12(1-4):187–92.

Obertreis B, Rutkowski T, et al. Ex-vivo in-vitro inhibition of lipopolysaccharide stimulated tumor necrosis factor-alpha and interleukin-1 beta secretion in human whole blood by extractum urticae dioicae foliorum. *Arzneimittelforschung.* 1996 May;46(4):389–94.

O'Leary KA, de Pascual-Tereasa S, et al. Effect of flavonoids and vitamin E on cyclooxygenase-2 (COX-2) transcription. *Mutat Res.* 2004 Jul;551(1-2):245–54.

Plummer SM, Holloway KA, et al. Inhibition of cyclo-oxygenase 2 expression in colon cells by the chemopreventive agent curcumin involves inhibition of NF-kappaB activation via the NIK/IKK signalling complex. *Oncogene.* 1999 Nov 28;18(44):6013–20.

Popoprigrorova VG, Bobylev AA, et al. [Peculiarities of free radical and antioxidant statute in patients with systemic lupus erythematosus and rheumatoid arthritis]. *Biomed Khim.* 2005 Sep-Oct;51(5):536–41.

Pullman-Mooar S, Laposata M, et al. Alteration of the cellular fatty acid profile and the production of eicosanoids in human monocytes by gamma-linolenic acid. *Arthritis Rheum.* 1990 Nov;33(10):1526–33.

Regan E, Flannelly J, et al. Extracellular superoxide dismutase and oxidant damage in osteoarthritis. *Arthritis Rheum.* 2005 Nov;52(11):3479–91.

Rovensky J, Kvetnansky R, et al. Hormone concentrations in synovial fluid of patients with rheumatoid arthritis. *Clin Exp Rheumatol.* 2005 May-Jun;23(3):292–6.

Safayhi H, Rall B, et al. Inhibition by boswellic acids of human leukocyte elastase. *J Pharmacol Exp Ther.* 1997 May;281(1):460–3.

Sarban S, Kocyigit A, et al. Plasma total antioxidant capacity, lipid peroxidation, and erythrocyte antioxidant enzyme activities in patients with rheumatoid arthritis and osteoarthritis. *Clin Biochem.* 2005 Nov;38(11):981–6.

Schmidt M, Weidler C, et al. Androgen conversion in osteoarthritis and rheumatoid arthritis synoviocytes: Androstenedione and testosterone inhibit estrogen formation and favor production of more potent 5alpha-reduced androgens. *Arthritis Res Ther.* 2005;7(5):R938–R948.

Schulze-Tanzil G, de SP, et al. Effects of the antirheumatic remedy hox alpha—a new stinging nettle leaf extract—on matrix metalloproteinases in human chondrocytes in vitro. *Histol Histopathol.* 2002 Apr;17(2):477–85.

Srivastava KC, Mustafa T. Ginger (*Zingiber officinale*) in rheumatism and musculoskeletal disorders. *Med Hypotheses*. 1992;39(4):342–8.

Straub RH, Harle P, et al. Sex hormone concentrations in patients with rheumatoid arthritis are not normalized during 12 weeks of anti-tumor necrosis factor therapy. *J Rheumatol*. 2005 Jul;32(7):1253–8.

Sundrarjun T, Komindr S, et al. Effects of n-3 fatty acids on serum interleukin-6, tumour necrosis factor-alpha and soluble tumour necrosis factor receptor p55 in active rheumatoid arthritis. *J Int Med Res*. 2004 Sep-Oct;32(5):443–54.

Tanaka S, Sato T, et al. Prevention of UVB-induced photoinflammation and photoaging by a polymethoxy flavonoid, nobiletin, in human keratinocytes in vivo and in vitro. *Biochem Pharmacol*. 2004 Aug 1;68(3):433–9.

Watson J, Byars ML, et al. Cytokine and prostaglandin production by monocytes of volunteers and rheumatoid arthritis patients treated with dietary supplements of blackcurrant seed oil. *Br J Rheumatol*. 1993;32(12):1055–8.

Weidler C, Struharova S, et al. Tumor necrosis factor inhibits conversion of dehydroepiandrosterone sulfate (DHEAS) to DHEA in rheumatoid arthritis synovial cells: A prerequisite for local androgen deficiency. *Arthritis Rheum*. 2005 Jun;52(6):1721–9.

All Contents Copyright © 1995-2009 Life Extension Foundation All rights reserved.

LifeExtension®

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.