

Low Dose Aspirin (Enteric Coated)

81 mg, 500 tablets

Item Catalog Number: 00892

Numerous studies document the multiple health benefits of daily low dose aspirin. Aspirin helps to maintain normal platelet aggregation in blood vessels via several mechanisms, including suppressing thromboxane H₂, prostaglandin E₂, and possibly C-reactive protein.¹¹⁶⁻¹³³

References

Active Ingredient (in each tablet)

Serving Size 1 tablet

Servings Per Container 500

Amount Per Serving

Aspirin

81 mg

Inactive ingredients: acetylated monoglycerides, anhydrous lactose, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10, hypromellose, hypromellose phthalate, iron oxide ochre, microcrystalline cellulose, mineral oil, polyethylene glycol (PEG-400), polysorbate 80, titanium dioxide.

Dosage and Use

- For the temporary relief of minor aches and pains or as recommended by your doctor.
- Adults and children 12 years and over: take four to eight tablets every four hours, not to exceed 48 tablets in 24 hours unless directed by a doctor.
- Drink a full glass of water with each dose.
- Children under 12 years: consult a doctor.

Caution

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Allergy alert

Aspirin may cause a severe allergic reaction which may include: hives, facial swelling, shock, asthma (wheezing).

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Alcohol warning: If you consume three or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding.

Ask a doctor or pharmacist before use if you are taking a prescription drug for anticoagulation (blood thinning), gout, diabetes, or arthritis.

Stop use and ask a doctor if new symptoms occur • an allergic reaction occurs. Seek medical help right away • pain gets worse or lasts more than 10 days • redness or swelling is present • ringing in the ears or loss of hearing occurs.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last three months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Ask a doctor before use if you have: asthma • ulcers • bleeding problems • stomach problems (such as heart-burn, upset stomach or stomach pain) that persist or recur.

Important

See your doctor before taking this product for other serious side effects that could occur with self treatment.

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.