

## Homocysteine Reduction

### REFERENCES

- Al-Obaidi MK, Stubbs PJ, et al. Elevated homocysteine levels are associated with increased ischemic myocardial injury in acute coronary syndromes. *J Am Coll Cardiol*. 2000 Oct;36(4):1217-1222.
- Anderson JL, Jensen KR, et al. Effect of folic acid fortification of food on homocysteine-related mortality. *Am J Med*. 2004 Feb 1;116(3):158-164.
- Anderson JL, Muhlestein JB, et al. Plasma homocysteine predicts mortality independently of traditional risk factors and C-reactive protein in patients with angiographically defined coronary artery disease. *Circulation*. 2000 Sep 12;102(11):1227-1232.
- Carlsen SM, Folling I, et al. Metformin increases total serum homocysteine levels in non-diabetic male patients with coronary heart disease. *Scand J Clin Lab Invest*. 1997 Oct;57(6):521-527.
- Charlton CG, Mack J. Substantia nigra degeneration and tyrosine hydroxylase depletion caused by excess S-adenosylmethionine in the rat brain: support for an excess methylation hypothesis for parkinsonism. *Mol Neurobiol*. 1994 Aug-Dec; 9(1-3):149-161.
- De Bree A, Verschuren WM, et al. Homocysteine determinants and the evidence to what extent homocysteine determines the risk of coronary heart disease. *Pharmacol Rev*. 2002 Dec;54(4):599-618.
- de la Vega MJ, Santolaria F, et al. High prevalence of hyperhomocysteinemia in chronic alcoholism: the importance of the thermolabile form of the enzyme methylenetetrahydrofolate reductase (MTHFR). *Alcohol*. 2001 Oct;25(2):59-67.
- Devlin TM. *Textbook of Biochemistry with Clinical Correlations*. 5th ed. New York, NY : Wiley-Liss; 2001.
- Dierkes J, Westphal S, et al. Serum homocysteine increases after therapy with fenofibrate or bezafibrate. *Lancet*. 1999 Jul 17;354(9174):219-220.
- Dudman NP, Guo XW, et al. Human homocysteine catabolism: three major pathways and their relevance to development of arterial occlusive disease. *J Nutr*. 1996 Apr;126(suppl 4):1295S-1300S.
- Fava M, Borus JS, et al. Folate, vitamin B12, and homocysteine in major depressive disorder. *Am J Psychiatry*. 1997 Mar;154(3):426-428.
- Finkelstein JD, Martin JJ. Methionine metabolism in mammals: distribution of homocysteine between competing pathways. *J Biol Chem*. 1984 Aug 10;259(15):9508-9513.
- Friedman AN, Bostom AG, et al. The kidney and homocysteine metabolism. *J Am Soc Nephrol*. 2001 Oct;12(10):2181-2189.
- Garg R, Malinow M, et al. Niacin treatment increases plasma homocyst(e)ine levels. *Am Heart J*. 1999 Dec;138(6 pt 1):1082-1087.
- Goren JL, Stoll AL, et al. Bioavailability and lack of toxicity of S-adenosyl-L-methionine (SAME) in humans. *Pharmacotherapy*. 2004 Nov;24(11):1501-1507.
- Haagsma CJ, Blom HJ, et al. Influence of sulfasalazine, methotrexate, and the combination of both on plasma homocysteine concentrations in patients with rheumatoid arthritis. *Ann Rheum Dis*. 1999 Feb;58(2):79-84.
- Harker LA, Ross R, et al. Homocystine-induced arteriosclerosis: the role of endothelial cell injury and platelet response in its genesis. *J Clin Invest*. 1976 Sep;58(3):731-741.
- Hvas AM, Juul S, et al. Vitamin B6 level is associated with symptoms of depression. *Psychother Psychosom*. 2004 Nov-Dec;73(6):340-343.

Jacques PF, Selhub J, et al. The effect of folic acid fortification on plasma folate and total homocysteine concentrations. *N Engl J Med*. 1999 May 13;340(19):1449-1454.

Jakubowski H. Homocysteine-thiolactone and S-nitroso-homocysteine mediate incorporation of homocysteine into protein in humans. *Clin Chem Lab Med*. 2003 Nov;41(11):1462-1466.

Joosten E, Lesaffre E, et al. Is metabolic evidence for vitamin B-12 and folate deficiency more frequent in elderly patients with Alzheimer's disease? *J Gerontol A Biol Sci Med Sci*. 1997 Mar;52(2):M76-M79.

Lamango NS, Charlton CG. Farnesyl-L-cysteine analogs block SAM-induced Parkinson's disease-like symptoms in rats. *Pharmacol Biochem Behav*. 2000 Aug;66(4):841-849.

Lee H, Kim JH, et al. Creatine synthesis and transport systems in the male rat reproductive tract. *Biol Reprod*. 1998 Jun;58(6):1437-1444.

Life Extension Foundation. *Anti-Aging News*. Hollywood, Fla : Life Extension Foundation, November 1981;85-86.

Lombardo F, Sgro P, et al. Might erectile dysfunction be due to the thermolabile variant of methylenetetrahydrofolate reductase? *J Endocrinol Invest*. 2004 Oct;27(9):883-885.

Malinow MR, Bostom AG, et al. Homocyst(e)ine, diet, and cardiovascular diseases: a statement for healthcare professionals from the Nutrition Committee, American Heart Association. *Circulation*. 1999 Jan 5-12;99(1):178-182.

Martinez-Chantar ML, Garcia-Trevijano ER, et al. Importance of a deficiency in S-adenosyl-L-methionine synthesis in the pathogenesis of liver injury. *Am J Clin Nutr*. 2002 Nov;76(5):1177S-1182S.

Matetzky S, Freimark D, et al. Association of elevated homocysteine levels with a higher risk of recurrent coronary events and mortality in patients with acute myocardial infarction. *Arch Intern Med*. 2003 Sep 8;163(16):1933-1937.

McCaddon A, Davies G, et al. Total serum homocysteine in senile dementia of Alzheimer type. *Int J Geriatr Psychiatry*. 1998 Apr;13(4):235-239.

McCully KS. Homocysteine and vascular disease. *Nat Med*. 1996 Apr;2(4):386-389.

McKay DL, Perrone G, et al. Multivitamin/mineral supplementation improves plasma B-vitamin status and homocysteine concentration in healthy older adults consuming a folate-fortified diet. *J Nutr*. 2000 Dec;130(12):3090-3096.

Morris MS, Bostom AG, et al. Hyperhomocysteinemia and hypercholesterolemia associated with hypothyroidism in the third US National Health and Nutrition Examination Survey. *Atherosclerosis*. 2001 Mar;155(1):195-200.

Muller T, Werne B, et al. Nigral endothelial dysfunction, homocysteine, and Parkinson's disease. *Lancet*. 1999 Jul 10;345(9173):126-127.

Neal B, MacMahon S, et al. Dose-dependent effects of folic acid on plasma homocysteine in a randomized trial conducted among 723 individuals with coronary heart disease. *Eur Heart J*. 2002 Oct;23(19):1509-1515.

Nygard O, Nordrehaug JE, et al. Plasma homocysteine levels and mortality in patients with coronary artery disease. *N Engl J Med*. 1997 Jul 24;337(4):230-236.

Oakley GP Jr. Eat right and take a multivitamin. *N Engl J Med*. 1998 Apr 9;338(15):1060-1061.

Ramos MI, Allen LH, et al. Plasma folate concentrations are associated with depressive symptoms in elderly Latina women despite folic acid fortification. *Am J Clin Nutr*. 2004 Oct;80(4):1024-1028.

Righetti M, Tommasi A, et al. Effective homocysteine-lowering vitamin B treatment in peritoneal dialysis patients. *Perit Dial Int*. 2004 Jul-Aug;24(4):373-377.

Schnyder G, Flammer Y, et al. Plasma homocysteine levels and late outcome after coronary angioplasty. *J Am Coll Cardiol*. 2002 Nov 20;40(10):1769-1776.

Schwanninger MC, Ringleb P, et al. Elevated plasma concentrations of homocysteine in antiepileptic drug treatment. *Epilepsia*. 1999 Mar;40(3):345-350.

Silveri MM, Parow AM, et al. S-adenosyl-L-methionine: effects on brain bioenergetic status and transverse relaxation time in healthy subjects. *Biol Psychiatry*. 2003 Oct 15;54(8):833-839.

Spence JD, Howard VJ, et al. Vitamin Intervention for Stroke Prevention (VISP) trial: rationale and design. *Neuroepidemiology*. 2001 Feb;20(1):16-25.

Stampfer MJ, Malinow MR, et al. A prospective study of plasma homocyst(e)ine and risk of myocardial infarction in US physicians. *JAMA*. 1992 Aug 19;268(7):877-881.

Stead LM, Au KP, et al. Methylation demand and homocysteine metabolism: effects of dietary provision of creatine and guanidinoacetate. *Am J Physiol Endocrinol Metab*. 2001 Nov;281(5):E1095-E1100.

Stuhlinger MC, Tsao PS, et al. Homocysteine impairs the nitric oxide synthase pathway: role of asymmetric dimethylarginine. *Circulation*. 2001 Nov 20;104(21): 2569-2575.

Tucker KL, Mahnken B, et al. Folic acid fortification of the food supply: potential benefits and risks for the elderly population. *JAMA*. 1996 Dec 18;276(23):1879-1885.

Vasan RS, Beiser A, et al. Plasma homocysteine and risk for congestive heart failure in adults without prior myocardial infarction. *JAMA*. 2003 Mar 12;289(10):1251-1257.

Venn BJ, Mann JI, et al. Dietary counseling to increase natural folate intake: a randomized, placebo-controlled trial in free-living subjects to assess effects on serum folate and plasma total homocysteine. *Am J Clin Nutr*. 2002 Oct;76(4):758-765.

Ventura P, Rosa MC, et al. Hyperhomocysteinaemia in chronic liver diseases: role of disease stage, vitamin status and methylenetetrahydrofolate reductase genetics. *Liver Int*. 2005 Feb;25(1):49-56.

Verhoef P, Stampfer MJ, et al. Homocysteine metabolism and risk of myocardial infarction: relation with vitamins B6, B12, and folate. *Am J Epidemiol*. 1996 May 1;143(9):845-859.

Vermeer SE, Hollander M, et al. Silent brain infarcts and white matter lesions increase stroke risk in the general population: the Rotterdam Scan study. *Stroke*. 2003 May;34(5):1126-1129.

Vine AK . Hyperhomocysteinemia: a risk factor for central retinal vein occlusion. *Am J Ophthalmol*. 2000 May;129(5):640-644.

All Contents Copyright © 1995-2009 Life Extension Foundation All rights reserved.

**LifeExtension®**

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.