

Thyroid Regulation

REFERENCES

- Althaus B, Staub JJ, et al. LDL/HDL-changes in subclinical hypothyroidism: Possible risk factors for coronary heart disease. *Clin Endocrinol (Oxf)*. 1988 Feb;28(2):157–63.
- Arnow WS. The heart and thyroid disease. *Clin Geriatr Med*. 1995;11:219–29.
- Arthur JR, Beckett GJ. Thyroid function. *Br Med Bull*. 1999;55(3):658–68.
- Bell DS, Ovalle F. Use of soy protein supplement and resultant need for increased dose of levothyroxine. *Endocr Pract*. 2001;7:193–4.
- Bellisola G, Bratter P, et al. The TSH-dependent variation of the essential elements iodine, selenium and zinc within human thyroid tissues. *J Trace Elem Med Biol*. 1998 Nov;12(3):177–82.
- Bunevicius R, Kazanavicius G, et al. Effects of thyroxine as compared with thyroxine plus triiodothyronine in patients with hypothyroidism. *N Engl J Med*. 1999 Feb 11;340(6):424–9.
- Danzi S, Klein I. Thyroid hormone and the cardiovascular system. *Minerva Endocrinol*. 2004 Sep;29(3):139–50.
- Dayan CM, Saravanan P, et al. Whose normal thyroid function is better—yours or mine? *Lancet*. 2002 Aug 3;360(9330):353.
- Delange F. Risks and benefits of iodine supplementation. *Lancet*. 1998 Mar 28;351(9107):923–4.
- Deshpande UR, Joseph LJ, et al. Effect of antioxidants (vitamin C, E and turmeric extract) on methimazole induced hypothyroidism in rats. *Indian J Exp Biol*. 2002 Jun;40(6):735–8.
- Duntas LH. Thyroid disease and lipids. *Thyroid*. 2002 Apr;12(4):287–93.
- Escobar-Morreale HF, Obregon MJ, et al. Replacement therapy for hypothyroidism with thyroxine alone does not ensure euthyroidism in all tissues as studied in thyroidectomized rats. *J Clin Invest*. 1995 Dec;96(6):2828–38.
- Fowler PB, McIvor J, et al. The effect of long-term thyroxine on bone mineral density and serum cholesterol. *J R Coll Physicians Lond*. 1996 Nov-Dec;30(6):527–32.
- Gaby AR. “Sub-laboratory” hypothyroidism and the empirical use of Armour thyroid. *Alt Med Rev*. 2004 Jun;9(2):157–79.
- Gartner R, Gasnier BC, et al. Selenium supplementation in patients with autoimmune thyroiditis decreases thyroid peroxidase antibodies concentrations. *J Clin Endocrinol Metab*. 2002 Apr;87(4):1687–91.
- Hak AE, Pols HA, et al. Subclinical hypothyroidism is an independent risk factor for atherosclerosis and myocardial infarction in elderly women: The Rotterdam Study. *Ann Intern Med*. 2000 Feb 15;132(4):270–8.
- Hennessey JV, Evaul JE, et al. L-thyroxine dosage: A reevaluation of therapy with contemporary preparations. *Ann Intern Med*. 1986 Jul;105(1):11–5.
- Hollowell JG, Staehling NW, et al. Serum TSH, T(4), and thyroid antibodies in the United States population (1988 to 1994): National Health and Nutrition Examination Survey (NHANES III). *J Clin Endocrinol Metab*. 2002 Feb;87(2):489–99.
- Jabbar MA, Larrea J, et al. Abnormal thyroid function tests in infants with congenital hypothyroidism: The influence of soy-based formula. *J Am Coll Nutr*. 1997 Jun;16(3):280–2.
- Jordan RM. Myxedema coma: Pathophysiology, therapy, and factors affecting prognosis. *Med Clin North Am*. 1995 Jan;79(1):185–

- Joshi JV, Bhandarkar SD, et al. Menstrual irregularities and lactation failure may precede thyroid dysfunction or goitre. *J Postgrad Med.* 1993;39:137–41.
- Kohrle J. The trace element selenium and the thyroid gland. *Biochimie.* 1999 May;81(5):527–33.
- Kralik A, Eder K, et al. Influence of zinc and selenium deficiency on parameters relating to thyroid hormone metabolism. *Horm Metab Res.* 1996 May;28(5):223–6.
- Kvetny J, Heldgaard PE, et al. Sub-clinical hypothyroidism is associated with a low-grade inflammation, increased triglyceride levels and predicts cardiovascular disease in males below 50 years. *Clin Endocrinol (Oxf).* 2004 Aug;61(2):232–8.
- Lorini R, Gastaldi R, et al. Hashimoto's thyroiditis. *Pediatr Endocrinol Rev.* 2003 Dec;1 Suppl 2:205–11; discussion 211.
- Nishiyama S, Futagoishi-Suginohara Y, et al. Zinc supplementation alters thyroid hormone metabolism in disabled patients with zinc deficiency. *J Am Coll Nutr.* 1994 Feb;13(1):62–7.
- O'Brien T, Dinneen SF, et al. Hyperlipidemia in patients with primary and secondary hypothyroidism. *Mayo Clin Proc.* 1993 Sep;68(9):860–6.
- Oner J, Kukner A, et al. Effect of vitamin E on follicular cell proliferation and expression of apoptosis-associated factors in rats with 6-N-propyl-2-thiouracil-induced goitrogenesis. *Folia Histochem Cytobiol.* 2003;41(4):213–7.
- Perk M, O'Neill BJ. The effect of thyroid hormone therapy on angiographic coronary artery disease progression. *Can J Cardiol.* 1997 Mar;13(3):273–6.
- Pollock MA, Sturrock A, et al. Thyroxine treatment in patients with symptoms of hypothyroidism but thyroid function tests within the reference range: Randomised double blind placebo controlled crossover trial. *BMJ.* 2001;323:891–5.
- Remer T, Neubert A, et al. Increased risk of iodine deficiency with vegetarian nutrition. *Br J Nutr.* 1999 Jan;81(1):45–9.
- Romaldini JH, Sgarbi JA, et al. [Subclinical thyroid disease: Subclinical hypothyroidism and hyperthyroidism]. *Arq Bras Endocrinol Metabol.* 2004 Feb;48(1):147–58.
- Ruz M, Codoceo J, et al. Single and multiple selenium-zinc-iodine deficiencies affect rat thyroid metabolism and ultrastructure. *J Nutr.* 1999 Jan;129(1):174–80.
- Saddock BJ, Saddock V. Psychological factors affecting medical conditions. In: Stoudeomire A, McDaniel JS, eds. *Kaplan and Saddocks Comprehensive Textbook of Psychiatry.* Philadelphia, Pa: Lippincott Williams and Willens; 2000:(vol 2)1765–888.
- Saito I, Ito K, et al. Hypothyroidism as a cause of hypertension. *Hypertension.* 1983 Jan-Feb;5(1):112–5.
- Sarandol E, Tas S, et al. Oxidative stress and serum paraoxonase activity in experimental hypothyroidism: Effect of vitamin E supplementation. *Cell Biochem Funct.* 2005 Jan-Feb;23(1):1–8.
- Sawaka AM, Gerstein HC, et al. Does a combination regimen of thyroxine (T4) and 3,5,3'-triiodothyronine improve depressive symptoms better than T4 alone in patients with hypothyroidism? Results of a double-blind randomized, controlled trial. *J Clin Endocrinol Metab.* 2003 Oct;88(10):4551–5.
- Smallridge RC. Metabolic and anatomic thyroid emergencies: A review. *Crit Care Med.* 1992 Feb;20(2):276–91.
- Stewart JC, Vidor GI. Thyrotoxicosis induced by iodine contamination of food: A common unrecognized condition? *Br Med J.* 1976 Feb 14;1(6006):372–5.
- Surks MI, Chopra IJ, et al. American Thyroid Association guidelines for use of laboratory tests in thyroid disorders. *JAMA.* 1990 Mar 16;263(11):1529–32.
- Tagawa N, Tamanaka J, et al. Serum dehydroepiandrosterone, dehydroepiandrosterone sulfate, and pregnenolone sulfate concentrations in patients with hyperthyroidism and hypothyroidism. *Clin Chem.* 2000;46:523–8.

Walsh JP, Shiels L, et al. Combined thyroxine/liothyronine treatment does not improve well-being, quality of life, or cognitive function compared to thyroxine alone: A randomized controlled trial in patients with primary hypothyroidism. *J Clin Endocrinol Metab.* 2003 Oct;88(10):4543–50.

Walsh JP, Stuckey BG. What is the optimal treatment for hypothyroidism? *Med J Aust.* 2001;174:41–3.

Westphal SA. Unusual presentations of hypothyroidism. *Am J Med Sci.* 1997 Nov;314(5):333–7.

Wortsman J, Rosner W, et al. Abnormal testicular function in men with primary hypothyroidism. *Am J Med.* 1987 Feb;82(2):207–12.

Zimmerman MB, Kohrle J. The impact of iron and selenium deficiencies on iodine and thyroid metabolism: Biochemistry and relevance to public health. *Thyroid.* 2002 Oct;12(10):867–78.

All Contents Copyright © 1995-2009 Life Extension Foundation All rights reserved.

LifeExtension®

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.