

Acne

REFERENCES

- Adebamowo CA, Spiegelman D et al. High school dietary dairy intake and teenage acne. *J Am Acad Dermatol*. 2005 Feb;52(2):207-14.
- Bassett IB, Pannowitz DL et al. A comparative study of tea-tree oil versus benzoyl peroxide in the treatment of acne. *Med J Austral* 1990;153:455-58.
- Bataille V, Snieder H et al. The influence of genetics and environmental factors in the pathogenesis of acne: a twin study of acne in women. *J Invest Dermatol* 2002 Dec; 119(6): 1317-22.
- Brown DJ, Dattner AM. Phytotherapeutic approaches to common dermatologic conditions. *Arch Dermatol* 1998; 134:1401-4.
- Carmina E, Lobo RA. A comparison of the relative efficacy of antiandrogens for the treatment of acne in hyperandrogenic women. *Clin Endocrinol (Oxf)*. 2002 Aug;57(2):231-34.
- Cordain L, Lindeberg S et al. Acne vulgaris: a disease of Western civilization. *Arch Dermatol*. 2002 Dec; 138(12):1584-90.
- Cunliffe WJ, Fernandez C et al. An observer-blind parallel-group, randomized, multicentre clinical and microbiological study of a topical clindamycin/zinc gel and a topical clindamycin lotion in patients with mild/moderate acne. *J Dermatolog Treat*. 2005;16(4):213-8.
- Draelos ZD, Matsubara A et al. The effect of 2 percent niacinamide on facial sebum production. *J Cosmet Laser Ther*. 2006 Jun;8(2):96-101.
- Dreno B, Foulc P et al. Effect of zinc gluconate on propionibacterium acnes resistance to erythromycin in patients with inflammatory acne: in vitro and in vivo study. *Eur J Dermatol* 2005 May-Jun; 15(3): 152-55.
- El-Akawi Z, Abdel-Latif N et al. Does the plasma level of vitamins A and E affect acne condition? *Clin Exp Dermatol* 2006 May; 31(3): 430-34.
- Elman M, Lask G. The role of pulsed light and heat energy (LHE) in acne clearance. *J Cosmet Laser Ther* 2004 Jun; 6(2): 91-95.
- Guenther LC. Optimizing treatment with topical tazarotene. *Am J Clin Dermatol*. 2003;4(3):197-202.
- Inui S, Nakajima T et al. Potential anti-androgenic activity of roxithromycin in skin. *J Dermatol Sci*. 2001 Oct;27(2):147-51.
- Karlsson T, Vahlquist A et al. 13-cis-retinoic acid competitively inhibits 3 alpha-hydroxysteroid oxidation by retinol dehydrogenase RoDH-4: a mechanism for its anti-androgenic effects in sebaceous glands? *Biochem Biophys Res Commun*. 2003 Mar 28;303(1):273-8.
- Kovalev VM. Complex treatment of acne with lipoic acid. *Vrach Delo* 1981a Mar; (3): 108-13.
- Kovalev VM. Lipoic acid content in the blood and urine in acne and its use in the treatment of this disease. *Vestn Dermatol Venerol* 1981b Mar; (3): 65-67.
- Layton AM. Optimal management of acne to prevent scarring and psychological sequelae. *Am J Clin Dermatol*. 2001;2(3):135-41.
- Lemay A, Poulin Y et al. Oral contraceptives as anti-androgenic treatment of acne. A comparison of the relative efficacy of antiandrogens for the treatment of acne in hyperandrogenic women. Clinical application of 5alpha-reductase inhibitors. *J Obstet Gynaecol Can*. 2002 Jul;24(7):559-67.
- Nussbaumer P, Billich A. Steroid sulfatase inhibitors. *Med Res Rev*. 2004 Jul; 24(4):529-76.

Omi T, Bjerring P et al. 420 nm intense continuous light therapy for acne. *J Cosmet Laser Ther* 2004 Nov; 6(3): 156-62.

Papageorgiou PP, Chu AC. Chloroxylenol and zinc oxide containing cream (Nels cream) vs. 5 percent benzoyl peroxide cream in the treatment of acne vulgaris. A double-blind, randomized, controlled trial. *Clin Exp Dermatol*. 2000 Jan;25(1):16-20.

Paranjpe P, Kulkarni PH. Comparative efficacy of four Ayurvedic formulations in the treatment of acne vulgaris: a double-blind randomized placebo-controlled clinical evaluation. *J Ethnopharm*. 1995; 49:127-32.

Pierard-Franchimont C, Goffin V et al. A double-blind controlled evaluation of the sebosuppressive activity of topical erythromycin-zinc complex. *Eur J Clin Pharmacol* 1995; 49(1-2): 57-60.

Puglia C, Tropea S et al. In vitro percutaneous absorption studies and in vivo evaluation of anti-inflammatory activity of essential fatty acids (EFA) from fish oil extracts. *Int J Pharm*. 2005 Aug 11;299(1-2):41-8.

Raman A, Weir U et al. Antimicrobial effects of tea-tree oil and its major components on *Staphylococcus aureus*, *Staph. epidermidis* and *Propionibacterium acnes*. *Lett Appl Microbiol* 1995 Oct; 21(4): 242-45.

Reifen R. vitamin A as an anti-inflammatory agent. *Proc Nutr Soc* 2002 Aug; 6(3): 397-400.

Rossen MH, Roed-Petersen J. [Acne vulgaris]. *Ugeskr Laeger*. 1993 Mar 15;155(11):775-8.

Russell JJ. Topical therapy for acne. *Am Fam Physician*. 2000 Jan 15;61(2):357-66.

Santos MA, Belo VG et al. Effectiveness of photodynamic therapy with topical 5-aminolevulinic acid and intense pulsed light versus intense pulsed light alone in the treatment of acne vulgaris: comparative study. *Dermatol Surg* 2005 Aug; 31(8 Pt 1): 910-15.

Shahbakhti H, Watson RE et al. Influence of eicosapentaenoic acid, an omega-3 fatty acid, on ultraviolet-B generation of prostaglandin-E2 and proinflammatory cytokines interleukin-1 beta, tumor necrosis factor-alpha, interleukin-6 and interleukin-8 in human skin in vivo. *Photochem Photobiol*. 2004 Sep-Oct;80(2):231-5.

Shalita AR, Smith JG et al. Topical nicotinamide compared with clindamycin gel in the treatment of inflammatory acne vulgaris. *Int J Dermatol* 1995 Jun; 34(6): 434-37.

Smith R, Mann N et al. The effect of short-term altered macronutrient status on acne vulgaris and biochemical markers of insulin sensitivity. *Asia Pac J Clin Nutr*. 2004;13(Suppl):S67.

Smolinski KN, Yan AC et al. Acne update: 2004. *Curr Opin Pediatr*. 2004 Aug 8;16(4):385-91.

Tremblay JF, Sire DJ et al. Light-emitting diode 415 nm in the treatment of inflammatory acne: an open-label, multicenter, pilot investigation. *J Cosmet Laser Ther* 2006 Apr; 8(1): 31-33.

Van Hoogdalem EJ, Tarpstra IJ et al. Evaluation of the effect of zinc acetate on the stratum corneum penetration kinetics of erythromycin in healthy male volunteers. *Skin Pharmacol*. 1996;9(2):104-10.

Webster GF, Guenther L et al. A multicenter, double-blind, randomized comparison study of the efficacy and tolerability of once-daily tazarotene 0.1 percent gel and adapalene 0.1 percent gel for the treatment of facial acne vulgaris. *Cutis* 2002 Feb; 69(2 Suppl): 4-11.

Weiss JS, Shavin JS. Topical retinoid and antibiotic combination therapy for acne management. *J Drugs Dermatol*. 2004 Mar;3(2):146-54.

Yeung CK, Chan HH. Cutaneous adverse effects of lithium: epidemiology and management. *Am J Clin Dermatol*. 2004;5(1):3-8.

Zhanel GG, Homenuik K et al. The glycyclines: a comparative review with the tetracyclines. *Drugs*. 2004;64(1):63-88.

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.