

## Male Hormone Restoration

### REFERENCES

- Ahlbom E, Prins GS et al. Testosterone protects cerebellar granule cells from oxidative stress-induced cell death through a receptor mediated mechanism. *Brain Res.* 2001 Feb 23;892(2):255-62.
- Ahmed SR, Boucher AE et al. Transdermal testosterone therapy in the treatment of male hypogonadism. *J Clin Endocrinol Metab.* 1988 Mar;66(3):546-51.
- Ali H, Baig M et al. Relationship of serum and seminal plasma zinc levels and serum testosterone in oligospermic and azospermic infertile men. *J Coll Physicians Surg Pak.* 2005 Nov;15(11):671-3.
- Anderson RA, Bancroft J et al. The effects of exogenous testosterone on sexuality and mood of normal men. *J Clin Endocrinol Metab.* 1992 Dec;75(6):1503-7.
- Barrett-Connor E, Goodman-Gruen D et al. Endogenous sex hormones and cognitive function in older men. *J Clin Endocrinol Metab.* 1999a Oct;84(10):3681-5.
- Barrett-Connor E, Von Muhlen DG et al. Bioavailable testosterone and depressed mood in older men: the Rancho Bernardo Study. *J Clin Endocrinol Metab.* 1999b Feb;84(2):573-7.
- Bhasin S. Effects of testosterone administration on fat distribution, insulin sensitivity, and atherosclerosis progression. *Clin Infect Dis.* 2003;37(suppl 2):S142-S149.
- Boyanov MA, Boneva Z et al. Testosterone supplementation in men with type 2 diabetes, visceral obesity and partial androgen deficiency. *Aging Male.* 2003 Mar;6(1):1-7.
- Boyle P, Robertson C et al. Updated meta-analysis of clinical trials of *Serenoa repens* extract in the treatment of symptomatic benign prostatic hyperplasia. *BJU Int.* 2004 Apr;93(6):751-6.
- Carnahan RM, Perry PJ. Depression in aging men: the role of testosterone. *Drugs Aging.* 2004;21(6):361-76.
- Cavallini G, Caracciolo S et al. Carnitine versus androgen administration in the treatment of sexual dysfunction, depressed mood, and fatigue associated with male aging. *Urology* 2004 Apr;63(4):641-6.
- Cherrier MM, Asthana S et al. Testosterone supplementation improves spatial and verbal memory in healthy older men. *Neurology.* 2001 Jul 10;57(1):80-8.
- Cherrier MM, Plymate S et al. Relationship between testosterone supplementation and insulin-like growth factor-I levels and cognition in healthy older men. *Psychoneuroendocrinology.* 2004 Jan;29(1):65-82.
- Cohen PG. The hypogonadal-obesity cycle: role of aromatase in modulating the testosterone-estradiol shunt: a major factor in the genesis of morbid obesity. *Med Hypotheses.* 1999 Jan;52(1):49-51.
- Colantoni A, Emanuele MA et al. Hepatic estrogen receptors and alcohol intake. *Mol Cell Endocrinol.* 2002 Jul 31;193(1-2):101-4.
- Dambe JE, Bergh A et al. The acute effect of estrogens on testosterone production appears not to be mediated by testicular estrogen receptors. *Mol Cell Endocrinol.* 1983;31(1):105-16.
- Davidson JM, Kwan M, et al. Hormonal replacement and sexuality in men. *Clin Endocrinol Metab.* 1982 Nov;11(3):599-623.
- Debruyne F, Koch G et al. Comparison of a phytotherapeutic agent (Permixon) with an alpha-blocker (Tamsulosin) in the treatment of benign prostatic hyperplasia: a 1-year randomized international study. *Eur Urol.* 2002 May;41(5):497-506.

Dobrzycki S, Serwatka W et al. An assessment of correlations of testosterone sex hormone levels and the extensiveness of coronary heart disease and the ejection fraction of the left ventricle in males. *Med Invest*. 2003 Aug;50(3-4):162-9.

Dunning TL, Ward GM. Testosterone replacement therapy: perceptions of recipients and partners. *J Adv Nurs*. 2004 Sep;47(5):467-74.

Dzugan SA, Arnold SR. Hypercholesterolemia treatment: a new hypothesis or just an accident? *Med Hypotheses*. 2002 Dec;59(6):751-6.

Ebert T, Jockenhovel F et al. The current status of therapy for symptomatic late-onset hypogonadism with transdermal testosterone gel. *Eur Urol*. 2005 Feb;47(2):137-46.

Eng ET, Williams D et al. Anti-aromatase chemicals in red wine. *Ann N Y Acad Sci*. 2002 Jun;963:239-46.

Fiorelli G, Picariello L et al. Estrogen synthesis in human colon cancer epithelial cells. *J Steroid Biochem Mol Biol*. 1999 Dec 31;71(5-6):223-30.

Gambelunghe C, Rossi R et al. Effects of chrysin on urinary testosterone levels in human males. *J Med Food*. 2003 Winter;6(4):387-90.

Gann PH, Hennekens CH et al. A prospective study of plasma hormone levels, nonhormonal factors, and development of benign prostatic hyperplasia. *Prostate*. 1995 Jan;26(1):40-9.

Gansser D, Spiteller G. Plant constituents interfering with human sex hormone-binding globulin: evaluation of a test method and its application to *Urtica dioica* root extracts. *Z Naturforsch C*. 1995 Jan-Feb;50(1-2):98-104.

Gerber GS, Fitzpatrick JM. The role of a lipido-sterolic extract of *Serenoa repens* in the management of lower urinary tract symptoms associated with benign prostatic hyperplasia. *BJU Int*. 2004 Aug;94(3):338-44.

Giannakopoulos X, Baltogiannis D et al. The lipido-sterolic extract of *Serenoa repens* in the treatment of benign prostatic hyperplasia: a comparison of two dosage regimens. *Adv Ther*. 2002 Nov;19(6):285-96.

Goldstat R, Briganti E et al. Transdermal testosterone therapy improves well-being, mood, and sexual function in premenopausal women. *Menopause*. 2003 Sep-Oct;10(5):390-8.

Hak AE, Witteman JC et al. Low levels of endogenous androgens increase the risk of atherosclerosis in elderly men: the Rotterdam study. *J Clin Endocrinol Metab*. 2002 Aug;87(8):3632-9.

Hammond J, Le Q et al. Testosterone-mediated neuroprotection through the androgen receptor in human primary neurons. *J Neurochem*. 2001 Jun;77(5):1319-26.

Harman SM 2005. Testosterone in older men after the Institute of Medicine Report: where do we go from here? *Climacteric*. 2005 Jun;8(2):124-35.

He F, Feng L. Effects of some micronutrients on partial androgen deficiency in the aging male [in Chinese]. *Zhonghua Nan Ke Xue*. 2005 Oct;11(10):784-6.

Herbst KL, Bhasin S. Testosterone action on skeletal muscle. *Curr Opin Clin Nutr Metab Care*. 2004 May;7(3):271-7.

Hirano T, Homma M et al. Effects of stinging nettle root extracts and their steroidal components on the Na<sup>+</sup>,K<sup>+</sup>-ATPase of the benign prostatic hyperplasia. *Planta Med*. 1994 Feb;60(1):30-3.

Hogervorst E, Bandelow S et al. Low free testosterone is an independent risk factor for Alzheimer's disease. *Exp Gerontol*. 2004 Nov-Dec;39(11-12):1633-9.

Hryb DJ, Khan MS et al. The effect of extracts of the roots of the stinging nettle (*Urtica dioica*) on the interaction of SHBG with its receptor on human prostatic membranes. *Planta Med*. 1995 Feb; 61(1):31-2.

Janowsky JS, Chavez B et al. Sex steroids modify working memory. *J Cogn Neurosci*. 2000 May;12(3):407-14.

Janowsky JS, Oviatt SK et al. Testosterone influences spatial cognition in older men. *Behav Neurosci*. 1994 Apr;108(2):325-32.

Jeppesen LL, Jorgensen HS et al. Decreased serum testosterone in men with acute ischemic stroke. *Arterioscler Thromb Vasc Biol*. 1996 Jun;16(6):749-54.

Jiang H, Chen Z. Studies on sex hormone-binding globulin in the prostate [in Chinese]. *Zhonghua Nan Ke Xue*. 2004 Jun;10(6):443-4, 8.

Kaya O, Gokdemir K et al. Zinc supplementation in rats subjected to acute swimming exercise: its effect on testosterone levels and relation with lactate. *Neuro Endocrinol Lett*. 2006 Apr 25;27(1-2):267-70.

Kellis JT Jr, Vickery LE. Inhibition of human estrogen synthetase (aromatase) by flavones. *Science*. 1984 Sep 7;225(4666):1032-4.

Kilic M, Baltaci AK et al. The effect of exhaustion exercise on thyroid hormones and testosterone levels of elite athletes receiving oral zinc. *Neuro Endocrinol Lett*. 2006 Apr 25;27(1-2):247-52.

Killinger DW, Perel E et al. The relationship between aromatase activity and body fat distribution. *Steroids*. 1987 Jul-Sep;50(1-3):61-72.

King JA, De Oliveira WL et al. Deficits in testosterone facilitate enhanced fear response. *Psychoneuroendocrinology*. 2005 May;30(4):333-40.

Kley HK, Deselaers T et al. Enhanced conversion of androstenedione to estrogens in obese males. *J Clin Endocrinol Metab*. 1980a Nov;51(5):1128-32.

Kley HK, Edelman P et al. Relationship of plasma sex hormones to different parameters of obesity in male subjects. *Metabolism*. 1980b Oct;29(11):1041-5.

Krieg M, Nass R et al. Effect of aging on endogenous level of 5 alpha-dihydrotestosterone, testosterone, estradiol, and estrone in epithelium and stroma of normal and hyperplastic human prostate. *J Clin Endocrinol Metab*. 1993 Aug;77(2):375-81.

Kris-Etherton PM, Hecker KD et al. Bioactive compounds in foods: their role in the prevention of cardiovascular disease and cancer. *Am J Med*. 2002 Dec 30;113(suppl 9B):71S-88S.

Lichius JJ, Muth C. The inhibiting effects of *Urtica dioica* root extracts on experimentally induced prostatic hyperplasia in the mouse. *Planta Med*. 1997 Aug;63(4):307-10.

Lopatkin N, Sivkov A et al. Long-term efficacy and safety of a combination of sabal and urtica extract for lower urinary tract symptoms: a placebo-controlled, double-blind, multicenter trial. *World J Urol*. 2005 Jun;23(2):139-46.

Marin P, Holmang S et al. The effects of testosterone treatment on body composition and metabolism in middle-aged obese men. *Int J Obes Relat Metab Disord*. 1992b Dec;16(12):991-7.

Marin P, Krotkiewski M et al. Androgen treatment of middle-aged, obese men: effects on metabolism, muscle and adipose tissues. *Eur J Med*. 1992a Oct;1(6):329-36.

Misao R, Nakanishi Y et al. Levels of sex hormone-binding globulin and corticosteroid-binding globulin mRNAs in corpus luteum of human subjects: correlation with serum steroid hormone levels. *Gynecol Endocrinol*. 1999 Apr;13(2):82-8.

Moffat SD, Zonderman AB et al. Longitudinal assessment of serum free testosterone concentration predicts memory performance and cognitive status in elderly men. *J Clin Endocrinol Metab*. 2002 Nov;87(11):5001-7.

Moger WH. Direct effects of estrogens on the endocrine function of the mammalian testis. *Can J Physiol Pharmacol*. 1980 Sep;58(9):1011-22.

Morgentaler A. Testosterone replacement therapy and prostate risks: where's the beef? *Can J Urol*. 2006 Feb;13(suppl 1):40-3.

Morley JE. Testosterone and behavior. *Clin Geriatr Med*. 2003 Aug;19(3):605-16.

Nankin HR, Calkins JH. Decreased bioavailable testosterone in aging normal and impotent men. *J Clin Endocrinol Metab*. 1986

Dec;63(6):1418-20.

Okun MS, DeLong MR et al. Plasma testosterone levels in Alzheimer and Parkinson diseases. *Neurology*. 2004 Feb 10;62(3):411-3.

Orengo CA, Fullerton G et al. Male depression: a review of gender concerns and testosterone therapy. *Geriatrics*. 2004 Oct;59(10):24-30.

Pardridge WM. Serum bioavailability of sex steroid hormones. *Clin Endocrinol Metab*. 1986 May;15(2):259-78.

Phillips GB. Relationship between serum sex hormones and the glucose-insulin-lipid defect in men with obesity. *Metabolism*. 1993 Jan;42(1):116-20.

Purohit V. Can alcohol promote aromatization of androgens to estrogens? A review. *Alcohol*. 2000 Nov;22(3):123-7.

Rabkin JG, Wagner GJ et al. Testosterone therapy for human immunodeficiency virus-positive men with and without hypogonadism. *J Clin Psychopharmacol*. 1999 Feb;19(1):19-27.

Ready RE, Friedman J et al. Testosterone deficiency and apathy in Parkinson's disease: a pilot study. *J Neurol Neurosurg Psychiatry*. 2004 Sep;75(9):1323-6.

Rhoden EL, Morgentaler A. Risks of testosterone-replacement therapy and recommendations for monitoring. *N Engl J Med*. 2004 Jan 29;350(5):482-92.

Rucker D, Ezzat S et al. IGF-I and testosterone levels as predictors of bone mineral density in healthy, community-dwelling men. *Clin Endocrinol (Oxf)*. 2004 Apr;60(4):491-9.

Safarinejad MR. *Urtica dioica* for treatment of benign prostatic hyperplasia: a prospective, randomized, double-blind, placebo-controlled, crossover study. *J Herb Pharmacother*. 2005;5(4):1-11.

Salminen EK, Portin RI et al. Associations between serum testosterone fall and cognitive function in prostate cancer patients. *Clin Cancer Res*. 2004 Nov 15;10(22):7575-82.

Schaeffer EM, Walsh PC et al. Risks of testosterone replacement. *N Engl J Med*. 2004 May 6;350(19):2004-6.

Schottner M, Gansser D et al. Lignans from the roots of *Urtica dioica* and their metabolites bind to human sex hormone binding globulin (SHBG). *Planta Med*. 1997 Dec;63(6):529-32.

Schweiger U, Deuschle M et al. Testosterone, gonadotropin, and cortisol secretion in male patients with major depression. *Psychosom Med*. 1999 May-Jun;61(3):292-6.

Seidman SN, Walsh BT. Testosterone and depression in aging men. *Am J Geriatr Psychiatry*. 1999 Winter;7(1):18-33.

Severi G, Morris HA et al. Circulating steroid hormones and the risk of prostate cancer. *Cancer Epidemiol Biomarkers Prev*. 2006 Jan;15(1):86-91.

Shibata Y, Ito K et al. Changes in the endocrine environment of the human prostate transition zone with aging: simultaneous quantitative analysis of prostatic sex steroids and comparison with human prostatic histological composition. *Prostate*. 2000 Jan;42(1):45-55.

Shores MM, Mocerri VM et al. Low testosterone is associated with decreased function and increased mortality risk: a preliminary study of men in a geriatric rehabilitation unit. *J Am Geriatr Soc*. 2004 Dec;52(12):2077-81.

Steiner MS, Raghoebar S. Antiestrogens and selective estrogen receptor modulators reduce prostate cancer risk. *World J Urol*. 2003 May;21(1):31-6.

Talalay P, Fahey JW. Phytochemicals from cruciferous plants protect against cancer by modulating carcinogen metabolism. *J Nutr*. 2001 Nov;131(suppl 11):3027S-33S.

Tenover JL. Male hormone replacement therapy including "andropause." *Endocrinol Metab Clin N Am*. 1998;27(4):969-87.

Tok EC, Ertunc D et al. The effect of circulating androgens on bone mineral density in postmenopausal women. *Maturitas*. 2004 Jul 15;48(3):235-42.

Valimaki VV, Alftan H et al. Serum estradiol, testosterone, and sex hormone-binding globulin as regulators of peak bone mass and bone turnover rate in young Finnish men. *J Clin Endocrinol Metab*. 2004 Aug;89(8):3785-9.

van den Beld AW, de Jong FH et al. Measures of bioavailable serum testosterone and estradiol and their relationships with muscle strength, bone density, and body composition in elderly men. *J Clin Endocrinol Metab*. 2000 Sep;85(9):3276-82.

Van Look PF, Frolich M. Effects of ethinyloestradiol on plasma levels of pituitary gonadotrophins, testicular steroids and sex hormone binding globulin in normal men. *Clin Endocrinol*. 1981 Mar;14(3):237-43.

Walle UK, Galijatovic A et al. Transport of the flavonoid chrysin and its conjugated metabolites by the human intestinal cell line Caco-2. *Biochem Pharmacol*. 1999 Aug 1;58(3):431-8.

Waynberg J. Aphrodisiacs: contributions to the clinical evaluation of the traditional use of *Ptychopetalum*. Paper presented at: the First International Congress on Ethnopharmacology; June 1990; France.

Wilt T, Ishani A et al. *Serenoa repens* for benign prostatic hyperplasia. *Cochrane Database Syst Rev*. 2002;3:CD001423.

Wright JV, Lenard L. *Maximize Your Vitality and Potency*. Petaluma, Calif: Smart Publications; 1999.

Zhang Y, Champagne N et al. Estrogen and androgen protection of human neurons against intracellular amyloid beta1-42 toxicity through heat shock protein 70. *J Neurosci*. 2004 Jun 9;24(23):5315-21.

Zhao SP, Li XP. The association of low plasma testosterone level with coronary artery disease in Chinese men. *Int J Cardiol*. 1998 Jan 31;63(2):161-4.

Zmuda JM, Fahrenbach MC. The effect of testosterone aromatization on high-density lipoprotein cholesterol level and postheparin lipolytic activity. *Metabolism* 1993 Apr;42(4):446-50.

All Contents Copyright © 1995-2009 Life Extension Foundation All rights reserved.

**LifeExtension®**

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.