

Boswella Topical Cream

4 oz jar

Item Catalog Number: 00258

Topical application of boswella cream may provide localized relief from fluid imbalance.

References

Supplement Facts

Ingredients

Methyl salicylate	10%
Capsaicin	0.025%
Other ingredients: <i>Boswella serrata</i> , vitamins A, C, and E, aloe vera gel, liposome.	

Dosage and Use

- Adults and children ten years of age and older: Apply to affected area three to four times daily.
- Children under ten years of age: Do not use, consult a doctor.
- In order to be effective, this product must be applied three to four times daily. Application less frequently may not provide optimum relief. Initial relief is usually noted within one or two weeks. If condition does not improve in four weeks, discontinue use of this product.

Caution

For external use only. Avoid contact with the eyes. If you are under a doctor's care for any serious condition, consult your doctor before using this product. If prone to allergic reaction from aspirin, salicylates and/or capsaicin, consult a doctor before using. If condition worsens, or if symptoms persist for more than seven days or clear up and occur again within a few days, discontinue use of this product and consult a doctor. If redness is present, discontinue use and consult a doctor. Do not apply to wounds or damaged skin. Do not bandage tightly. Do not use with a heating pad (may blister skin).

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.