

Vinpocetine

5 mg, 100 tablets

Item Catalog Number: 00427

Vinpocetine is derived from vincamine, the major indole alkaloid of the periwinkle plant. No toxic effects have been seen from vinpocetine use at levels far above those recommended for this product.

Vinpocetine taken orally is easily absorbed.^{94,95} When taken orally, vinpocetine can:

- Improve blood supply to the brain⁹⁶⁻¹⁰¹
- Increase oxygen and glucose use by the brain^{96,102-104}
- Improve brain tolerance to hypoxic injury¹⁰⁵⁻¹⁰⁷
- Increase vasodilation response to hypoxia^{108,109}
- Maintain optimal energy of healthy brains^{96,102-104,110,111}
- Maintain normal coagulation of blood¹¹²
- Maintain healthy levels of some neurotransmitters^{97,113,114}
- Promote healthy attention, memory and concentration^{98,115-123}

References

Supplement Facts

Serving Size 2 tablets

Servings Per Container 50

Amount Per Serving

Vinpocetine

10 mg

Other ingredients: dicalcium phosphate, cellulose, stearic acid, magnesium stearate.

Contains soybeans. This product contains NO milk, egg, fish, peanuts, crustacean shellfish (lobster, crab, shrimp), tree nuts, wheat, yeast, gluten, corn, or rice. Contains NO sugar, and no artificial sweeteners, flavors, colors, or preservatives.

Dosage and Use

- Take two tablets once or twice daily with meals, or as recommended by a healthcare practitioner.

Caution

If you are taking warfarin (Coumadin®), consult your physician before using this product.

Warnings

- Keep out of reach of children.
- Do not exceed recommended dose.
- Do not purchase if outer seal is broken or damaged.
- If you have a bad reaction to product discontinue use immediately.
- When using nutritional supplements, please consult with your physician if you are undergoing treatment for a medical condition or if you are pregnant or lactating.

not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.