

## PSA Free/Total Ratio Reflex

This test is used to measure the percentage of free PSA relative to the total amount of PSA in men with serum PSA concentrations. Serum PSA levels increase due to physical changes to prostate architecture due to trauma, infection, inflammation and disease. Sensitivity of PSA levels to these changes serves as a basis for this test. Higher percentage of free PSA is associated with a lower risk of prostate cancer.

The blood test services are for informational purposes only. It is not the intention of National Diagnostics, Inc and Life Extension to provide specific medical advice but rather to provide users with information to better understand their health. Specific medical advice including diagnosis and treatment will not be provided. Always seek the advice of a trained health professional for medical advice, diagnosis or treatment.

Both the physician and the testing laboratory are independent contractors with whom National Diagnostics, Inc makes arrangements for your blood tests. Neither National Diagnostics, Inc nor Life Extension will be liable for any acts or commissions of the physician, the testing laboratory, or their agents or employees.

All Contents Copyright © 1995-2009 Life Extension Foundation All rights reserved.

**LifeExtension®**

These statements have not been evaluated by the FDA. These products are not intended to diagnose, treat, cure or prevent any disease. The information provided on this site is for informational purposes only and is not intended as a substitute for advice from your physician or other health care professional or any information contained on or in any product label or packaging. You should not use the information on this site for diagnosis or treatment of any health problem or for prescription of any medication or other treatment. You should consult with a healthcare professional before starting any diet, exercise or supplementation program, before taking any medication, or if you have or suspect you might have a health problem. You should not stop taking any medication without first consulting your physician.